

Neutrino and Parity Violation in Weak Interaction

Kiyoung Kim

Department of Physics, University of Utah, SLC, UT 84112

Abstract

It is shown that neutrino oscillation and Majorana type neutrino are not compatible with Special Theory of Relativity. Instead of the Majorana type neutrino, traditional neutrino ($m_\nu = 0$) is considered with additional assumptions that ν_R and $\bar{\nu}_L$ neutrinos exist in nature and naturally thus parity is conserved. Besides, neutrino itself is considered as *longitudinal vacuum-string oscillations*. To explain parity conservation $W^\pm$ bosons are suggested as momentary interacting states between $e^\pm$ and virtual-positive(negative)-charge strings, respectively. With these postulations, an alternative explanation is also suggested for solar neutrino problem.

1 Introduction

As the most amazing and even paradoxical one in elementary particles, neutrino has been known to us for a long time since 1931, when Wolfgang Pauli proposed a new particle – very penetrating and neutral – in β -decay. Nevertheless, we don't know much about neutrino itself. That is mainly because of experimental difficulty to detect neutrino itself and neutrino events too. Up to now, we are not quite sure if neutrino has a rest mass or not, and also we cannot explain *solar neutrino deficit*[25] [26][27][28] contrary to the expectation from Standard Solar Model[28]. Furthermore, Nature doesn't seem to be fair in *Weak Interaction* in which neutrino is involved. That has been known as *Parity Violation*.

As one of trial solutions to explain the *solar neutrino deficit*, neutrino *mass* or *flavor* oscillation has been suggested, and many experiments have been done and doing until now. However, in the theory of neutrino flavor oscillation, it is supposed that neutrino, for instance, ν_e , ν_μ , or ν_τ , is not in a fundamental state in Quantum mechanical point of view, but in a mixed state with *mass eigenstates*, such as m_1 , m_2 , and m_3 ($m_1 \neq m_2 \neq m_3$). This supposition implies that neutrino has an intrinsic structure and, thus, one of flavors spontaneously transforms to another kind of flavors, like the conversion of Meson K^0 into its antiparticle $\bar{K}^0$. If neutrino flavor mixing is true, we have another dilemma in *lepton number* conservation, which has been quite acceptable in phenomenology. In section(1.2), we are going to compare neutrino flavor mixing with the K^0 and $\bar{K}^0$ mixing. Moreover, the theory itself will be reviewed if it is physically sound or not.

In Weak Interaction, we know that *parity* is violated. When C. S. Wu *et al* showed parity violation in their experiment[1], it was astonishing because we had believed in a fairness in natural phenomena. Up to now, we don't know the reason for *parity violation*, and also we cannot explain why *parity violation* is detected only in Weak Interaction. Should we accept this fact as nature itself? Otherwise, there should be a reasonable reason.

As mentioned above, the experimentation is difficult because neutrinos have extremely small cross-sections(typical cross-section : 10^{-44} m^2 (1GeV)[2]); furthermore, neutrinos participate in only Weak Interaction in which the interaction range and the relative strength are $\ll 1\text{fm}(\sim 10^{-18}\text{m})$ and 10^{-15} , respectively.[4] Meanwhile, if the interaction range and the relative strength are compared with Electromagnetic Interaction case(range : ∞ , relative strength : 10^{-2})[4], it is not easy [for us] to understand why Electromagnetic Interaction - intermediated by *photon* - is suppressed in the Weak Interaction range despite that its interaction range is ∞ and the relative strength is 10^{-2} which is 10^{13} times bigger than Weak Interaction one.[3]

1.1 Fermion and Boson

Through the Complex Space model[5], it was possible to understand how Special Theory of Relativity and Quantum Mechanics can be connected, what is the physical entity of wave function - ontological point of view - in Quantum Mechanics, and how Schrödinger equation is related to the Complex Space model[6]. In that, Dirac's *hole theory*[7] was re-interpreted as following : In the Complex Space, vacuum particles - vacuum electrons in the model - are not packed completely; they can transfer energy(for instance, electromagnetic energy) through a wave motion such as *vacuum-string-oscillation*. In which the electromagnetic energy is propagated along the imaginary axis, and it is realized to real space through U(1) symmetry.

In Quantum Physics, we know that the *phase factor* of wave function is arbitrary if we consider only probability density($\psi \cdot \psi^*$); yet, it need to be considered in physical interactions. Especially,

its peculiarity was shown by Berry, M.V. in considering slowly evolving Quantum phase in the interaction of fermion(for example, electron) with magnetic field, and it has been known as *Berry phase*. [8][9]

Now, let us think about what the electron's spin is and, in general, what the fermion(electron, proton, neutron, etc) is. First of all, we know that the electron's spin can be represented as a rotation in *two dimensional complex space*, that is $SU(2)$; and the spin vector is realized through an interaction in which the interaction Hamiltonian is $\alpha(\vec{\sigma} \cdot \vec{r})(\sigma:\text{pauli matrix, } \alpha:\text{constant})$. [8] Furthermore, we can physically surmise that the peculiarity of Quantum phase occurs when the spin vector in $SU(2)$ is realized through the interaction with magnetic field because of the difference between $SU(2)$ and $SO(3)$ – two to one correspondence in their parameter spaces even though both are similar groups. Here, we can be sure, at least, that electron's spin vector(axial vector) resides in the Complex Space. According to the interpretation of Quantum formalism in




Figure 1: phase change through space inversion

the Complex Space [6], Quantum phase corresponds to the phase itself in the Complex Space. Although the Quantum phase in a stationary system is arbitrary – $U(1)$ – and doesn't affect physical interpretation, in a dynamical system the phase is no more free to choose. As a specific and simple example of Berry phase, let us imagine following : One electron (spin $\hbar/2$) is at origin under the influence of magnetic field ($\vec{H}$) which is on x - y plane with pointing to the origin. Now, let the field pointing direction rotate slowly around z axis; then, the system is in $SO(2) \subset SO(3)$ in real space and $U(1) \subset SU(2)$ in the Complex Space because of the correspondence between $SU(2)$ and $SO(3)$; $U(1)$ and $SO(2)$ in their transformations. Here, $U(1)$ is different from the one we traditionally use since our concern is $U(1)$ in the internal Complex Space. Now let us say, the electron's spin points to the positive x axis at the beginning. And let the field direction

rotate with ϕ degrees which corresponds to $\phi/2$ rotating simultaneously in the internal Complex Space. After a round trip of the system($\phi = 2\pi$) in real space, we can easily recognize that the spin points to the other direction – the negative x axis.

From Pauli exclusion principle we know that any two identical fermions cannot be at the same Quantum state, but bosons can share the same state. That means, the wave function representing two identical fermion system is anti-symmetry, but the wave function of two identical bosons is symmetry. To demonstrate this fact, let us say, $\Psi(1, 2)$ represents the Quantum state for two identical fermions. Then, $\Psi(2, 1) = -\Pi\Psi(1, 2) = -P\Psi(1, 2)$, where Π and P are respectively *exchange* and *parity* operator and it was used the fact that exchange operator(Π) is identical to parity operator(P) for this two identical fermion system. In Fig.(1) total spin vector(axial) in two identical fermion system is in internal Complex Space. After space inversion or parity operation the Quantum phase is changed from θ to $\pi + \theta$ with showing how the anti-symmetric wave function(two-fermion system) and the parity operation are connected.

With this fact we can distinguish *fermion* and *boson*. Fermion has a spin – intrinsic angular moment(half integer $\times \hbar$) – in Complex Space; boson can also have a spin, but the spin vector (axial) is in real space.

1.2 Neutrino Oscillation

One of long standing questions related elusive particle, neutrino, is neutrino oscillation. To understand the theory of neutrino oscillation, firstly we have to assume that each flavor of neutrinos(ν_e, ν_μ, ν_τ) has a rest mass. Of course, $m_{\nu_e} \neq m_{\nu_\mu} \neq m_{\nu_\tau}$ to be meaningful in the theory. Furthermore we have to suppose that there are fundamental Quantum states, in which the individual mass eigenstate doesn't appear in phenomenon, but linear combinations of these states emerge to ν_e, ν_μ or ν_τ .

In the respect of Quantum formalism : To describe *one* particle system, the wave function is a linear combination with all possible Quantum states – orthogonal and complete set of bases. There, each state represents *the particle* itself with a probability. Now, if the particle changes its identity, for instance, from ν_e to ν_μ , can we still use the same complete basis set to describe the new one? Or should we change the basis set to new one also? According to the theory, we can use the same basis set to describe these phenomenologically different particles because these particles are fundamentally identical but in different states to each other. However, K^0 and $\bar{K}^0$ mixing is different from neutrino flavor mixing case because they have internal structures and thus can be different intrinsic parities. In phenomenon, they appear to have different properties and transform spontaneously to each other with exchanging $\pi^\pm$ (Pions).

Now what about neutrinos? Do they have been known they have internal structures and different intrinsic parities? Although it has been known that neutrinos have imaginary intrinsic parities[12], it doesn't make any sense in phenomenology. What about in the middle of oscillation? For instance, "50% ν_e and 50% ν_μ " is possible? Although we can use Quantum formalism to describe a physical system, the particle's identity in the physical system should be clear at any time. Otherwise, there should be an additional explanation why the fundamental mass states are beyond physical phenomena.

Energy and Momentum conservation: Let us say, rest mass m_{ν_e} – electron neutrino – is moving in an isolated system with velocity β_e at $t = 0$, and after τ seconds(τ is arbitrary) the mass is changed to rest mass m_{ν_μ} ($m_{\nu_\mu} \neq m_{\nu_e}$) – muon neutrino – and the velocity to β_μ . First of all, energy conservation should be satisfied as

$$m_{\nu_e}\gamma_e = m_{\nu_\mu}\gamma_\mu \quad (1)$$

where $\gamma_e = 1/\sqrt{1 - \beta_e^2}$, $\gamma_\mu = 1/\sqrt{1 - \beta_\mu^2}$, and $c \equiv 1$, Moreover their momentums also should be conserved as

$$m_{\nu_e}\gamma_e\beta_e = m_{\nu_\mu}\gamma_\mu\beta_\mu \quad (2)$$

If the masses, m_{ν_e} and m_{ν_μ} are different, both equations, Eqn.(1) and Eqn.(2), cannot be satisfied simultaneously.

Even though our concerning is different in Special Theory of Relativity and Quantum Physics, both theories should be equally satisfied by a new theory because they are basic theories in physics and connected fundamentally. [5]

2 Dirac equation and Majorana neutrino

To describe spin $\frac{1}{2}\hbar$ particle(fermion), two formalisms – Dirac[13][14] and Majorana[10][11] [12] formalism – have been known. Since both formalisms are not derived directly from Schödinger equation – that is not deductive but inductive, it is natural [for us] to check if these two formalisms are physically enough meaningful or not.

In Majorana case; neutrino and antineutrino are identical such as $\nu_L \equiv \bar{\nu}_L$ and $\nu_R \equiv \bar{\nu}_R$ (Majorana abbreviation), and neutrino has a rest mass($m_\nu \neq 0$). Although the first condition is tolerable in the respect of phenomenological facts that we couldn't have detected $\bar{\nu}_L$ and ν_R , the second one is not compatible with Special Theory of Relativity.

If $m_\nu \neq 0$ no matter how small it is¹, then we can find the neutrino rest frame through proper Lorentz transformation; furthermore, we can even flip the helicity from right-handed($\bar{\nu}_R$) to left-handed(ν_L) or *vice versa*. Because there are only two kinds of neutrinos, ν_L and $\bar{\nu}_R$, the flipped one is always corresponded to the other one. However, obviously we can distinguish ν_L from $\bar{\nu}_R$ in their properties.[16][17] For instance, in interactions; $\nu_\mu + e \longrightarrow \nu_\mu + e$ and $\bar{\nu}_\mu + e \longrightarrow \bar{\nu}_\mu + e$, their total cross sections, $\sigma_{\nu e}$ and $\sigma_{\bar{\nu}e}$, are different.[17] If Majorana neutrinos can be applicable in physics, they should be identical in their physical properties, that is, $\nu_L \equiv \bar{\nu}_R$. Yet, the total cross section can be changed suddenly at a critical boost velocity in the transformation in spite of the fact that the cross section should be invariant. That means, a physical fact, which should be unique in all Lorentz frames, can be changed through the transformation. Therefore, both conditions in Majorana formalism cannot be feasible in physical situation as long as Special Theory of Relativity is impeccable. About the neutrino mass, there was already a similar argument in the respect of Group Theory in 1957[18]. Therefore, neutrinos satisfied by both conditions are not appropriate in physics.

If we abandon the first condition but still assume that neutrino mass is not zero, we have to treat neutrinos as like other spin $\frac{1}{2}\hbar$ fermions – not Majorana type fermions. However, what if we give up the mass of neutrino but hold the first condition, $\nu_L \equiv \bar{\nu}_L$ and $\nu_R \equiv \bar{\nu}_R$? Then, we have to ask again the old questions if ν_R and $\bar{\nu}_L$ exist or not; if these two neutrinos exist, why we cannot detect them; why parity is violated in Weak Interaction. As a possible case, let us assume that neutrino has no mass ($m_\nu = 0$) and that ν_R and $\bar{\nu}_L$ exist but we couldn't have detected them yet. If this supposition, which corresponds to Majorana neutrino with setting m_ν to zero[11], is true, we can understand why parity violation happens in Weak Interaction; moreover, we can find a clue to understand *solar neutrino deficit* problem.

In Dirac formalism; four spinors are closed by *parity* operation in general, and in neutrino case($m_\nu = 0$) the Dirac equation is decoupled to two component theory – Weyl equations. To compare spinor parts among neutrinos(ν_R , ν_L , $\bar{\nu}_R$, and $\bar{\nu}_L$) in free space, let us say :

$$\begin{aligned}\nu_R(x) &= \nu_R(k)e^{-ikx} & ; & \quad \nu_L(x) = \nu_L(k)e^{-ikx} \\ \bar{\nu}_R(x) &= \bar{\nu}_R(k)e^{ikx} & ; & \quad \bar{\nu}_L(x) = \bar{\nu}_L(k)e^{ikx}\end{aligned}\tag{3}$$

for a given k . Here, ν_R represents positive energy with right-handed neutrino; $\bar{\nu}_L$, negative energy with left-handed antineutrino, and *etc.* Using *charge conjugation* operator C which is consistent with the Dirac's hole interpretation, we can see that there are only two independent solutions[15] as

$$\begin{aligned}\bar{\nu}_R(k) &= C\gamma^0\nu_L^*(k) = -\nu_R(k) & \text{and} \\ \bar{\nu}_L(k) &= C\gamma^0\nu_R^*(k) = -\nu_L(k),\end{aligned}\tag{4}$$

¹our main concern here is about neutrino.

in which $\gamma^o = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$ and $C = \begin{pmatrix} 0 & -i\sigma_2 \\ -i\sigma_2 & 0 \end{pmatrix}$; the minus sign on RHS can be interpreted as the phase factor came from *Pauli exclusion* principle as in Sec.(1.1). With a normalization condition given for massless particles(neutrinos)[15], that is, $\nu_R(k)^\dagger \nu_R(k) = \nu_L(k)^\dagger \nu_L(k) = \bar{\nu}_R(k)^\dagger \bar{\nu}_R(k) = \bar{\nu}_L(k)^\dagger \bar{\nu}_L(k) = 2E_k$, plane wave solutions in Standard representation can be constructed as

$$\begin{aligned} \nu_R(k) &= \sqrt{E_k} \begin{pmatrix} +1 \\ 0 \\ +1 \\ 0 \end{pmatrix} & \bar{\nu}_R(k) &= \sqrt{E_k} \begin{pmatrix} -1 \\ 0 \\ -1 \\ 0 \end{pmatrix} \\ \nu_L(k) &= \sqrt{E_k} \begin{pmatrix} 0 \\ +1 \\ 0 \\ -1 \end{pmatrix} & \bar{\nu}_L(k) &= \sqrt{E_k} \begin{pmatrix} 0 \\ -1 \\ 0 \\ +1 \end{pmatrix} \end{aligned}$$

3 Neutrino

If neutrino mass is zero as assumed in the last Section, there are only two physically distinctive neutrinos for each kind, and their helicities are uniquely fixed to *right-handed* or *left-handed* once for all because of $m_\nu = 0$. Moreover, if we assume that ν_R and $\bar{\nu}_L$ exist, only the helicity of neutrino should be considered to distinguish neutrinos for each kind; and Dirac equation is closed by parity operation like other spin $\frac{1}{2}\hbar$ fermions. Phenomenological facts – neutrinos have an extremely small cross section, and Electromagnetic Interaction is highly suppressed in Weak Interaction limit($\sim 10^{-18}m$) – can be a good clue to understand neutrino. In what follows, we are going to propose what is *neutrino* and what is a possible model of $W^\pm$ boson with which *parity violation* can be explained.

3.1 Propagation mode

Since Electromagnetic wave, on which rest mass cannot be defined, was considered as *vacuum string oscillation*[5], we might consider a similarity of neutrino to photon. In the model of electromagnetic wave propagation[5] Planck's constant(h) is given as

$$h = 2\pi^2 m_e c \left(\frac{A^2}{d} \right), \quad (5)$$

where A is the *constant* amplitude for each wave string, d is the distance between two adjacent vacuum electrons, and m_e is bare electron mass. For a rough estimation if we use electron mass

instead of the bare mass, electron Compton wavelength($\lambda_c/2\pi$) has relation as following

$$\begin{aligned}\frac{S}{d} &\approx \frac{\hbar}{m_e c}, \\ &= 3.86 \times 10^{-13} m,\end{aligned}\tag{6}$$

in which $S = \pi A^2$. Now, if we accept A as $\sim 10^{-18} m$ ² because Electromagnetic Interaction is highly suppressed in this limit and, thus, we can surmise that electromagnetic wave cannot propagate in the limit. Then, d is estimated as $\sim 10^{-23} m$ from Eqn.(6). Furthermore, there is a limit of photon energy because d is not zero – the vacuum string oscillation is not through a continuous medium. For example, wavelengths of TeV and PeV energy photons³ are $\sim 10^{-18} m$ and $\sim 10^{-21} m$, respectively. But EeV energy photon is not possible because the wavelength($\lambda \sim 10^{-24} m$) is smaller than d . As a reference, PeV energy order γ rays have been detected in cosmic ray experiments.[19]

For neutrino to propagate Weak Interaction region($\sim 10^{-18} m$) there is a mode in which only two independent states exist, that is *longitudinal vacuum string oscillation* in which spin state(helicity) is also transferred through the string oscillation. With this model we can estimate bare electron size by a crude comparison of Compton scattering with neutrino and electron elastic scattering(ν_e, e). Let us say, the ratio of Compton scattering total cross section⁴ to the elastic scattering cross section $\sigma(\nu_e e)$ is

$$R = \frac{\sigma_{com}}{\sigma_{\nu e}} \sim \frac{A^2}{\delta^2},$$

where δ is bare electron radius. For 1 MeV of ν_e and photon in electron rest frame, $\sigma_{com} \sim 10^{-29} m^2$ and $\sigma_{\nu e} \sim 10^{-48} m^2$. [20] Then, the ratio, $R \sim 10^{19}$, and the bare electron radius can be estimated as $\delta \sim 10^{-27} m$.

3.2 $W^\pm$ boson and parity violation

Through experimental facts we have known that $\bar{\nu}_L$ and ν_R don't exist in nature. Yet we need to distinguish the existence in phenomenology and in ontology. If $\bar{\nu}_L$ and ν_R exist, necessarily we need an explanation why these two neutrinos couldn't have been detected. For example, in β^+ decay of ${}^8B \rightarrow {}^8Be^* + e^+ + \nu_e$ one proton is converted to a neutron inside the nucleus with emitting a positron e^+ and an electron neutrino ν_e . That is $p \rightarrow n + e^+ + \nu_e$. In this process of Weak Interaction it has been known that W^+ boson intermediate the process and the neutrino ν_e is *left handed* electron neutrino. However, if we assume W^+ boson($\Gamma_W^{e\nu} \simeq 0.23$ GeV, $\tau \sim 3 \times 10^{-24} sec$ [22]) as a momentary interacting state of a positron and *virtual positive*

²in facts, it can be even bigger than $\sim 10^{-18} m$.

³TeV = $10^{12} eV$, PeV = $10^{15} eV$, EeV = $10^{18} eV$

⁴low energy γ ray to avoid hadronic interaction

charge strings moving around the positron, we can figure out from Fig.(2) that there is no more preference in the neutrino helicity if the magnetic field is turned off.

In Fig.(2) e^+ represents bare charge of the positron, and q_v^+ stands for the virtual positive charge strings induced from vacuum polarization[21]. In which the virtual positive charge strings might even have a distribution depending on radial distance from the positron. Although we don't expect a positively induced virtual charges from the vacuum polarization if the positron is in a free space, inside Weak Interaction region($r \leq 10^{-18}m$) the virtual positive charges can be accumulated around the edge of the region and even experienced a repulsive force from the edge. According to the mirror image in Dirac's hole theory the virtual-positive-charges behave like that they have *positive* masses in phenomena. Since the state, interacting between the positron and




Figure 2: W^+ in β^+ decay

virtual-positive-charges, is not stable, soon it decay to a free positron and a neutrino, in which the kinetic energy of the virtual-positive-charges is transferred to the longitudinal vacuum string oscillation, that can be interpreted as neutrino as assumed before. Now, if magnetic field $\vec{H}$ is applied as in Fig.(2) to investigate the positron's spin orientation we can use Faraday Induction

law, which is an empirical and macroscopic law, because the radius of $10^{-18}m$ is much bigger than the bare electron size($10^{-27}m$). Then, the virtual-positive-charges rotating around the positron move to the negative z-axis(attractive in Faraday Induction law) and finally it will be transferred to the longitudinal vacuum string oscillation – neutrino – following negative imaginary z-axis as in the case of photon[5] because neutrino mass m_ν is assumed to be zero. On the other hand, the positron must move to the positive z-axis because of momentum conservation. If total spin of the system was $1\hbar$ pointing to the positive z-axis, the emitting electron neutrino definitely has *left handed* helicity. What if the magnetic fields is turned off? we cannot say which direction the neutrino choose for its emitting. From a reasonable guess, we can say that it should be equally probable for each direction.

In β^- decay, that is $n \rightarrow p + e^- + \nu_e$, W^- boson also can be treated similarly as in W^+ boson case if we remind of the mirror image in Dirac's hole theory, where W^- boson is considered as a momentary state of an electron e^- and virtual-negative-charge strings moving around the electron, and the virtual-negative-charge strings behave like that they have negative masses. With a similar set up like Fig.(2) except negative charges and the other rotating direction of the virtual-negative- charge strings, we can find out that the emitting electron neutrino is now *right handed* if the magnetic fields is on, and equally probable to be *right handed* or *left handed* if the magnetic field is off. Similarly, we can assume further that Z^0 boson is a momentary interacting state($\sim 3 \times 10^{-24}sec$) between virtual-positive-charge strings and virtual-negative-charge strings inside the Weak Interaction region($r \leq 10^{-18}m$).

From the reasoning as above, it is reasonable to say that the *parity violation* can be happened if we consider only one part of phenomena, in which we can investigate the spin orientation of lepton using magnetic field; however, *intrinsically* parity is conserved if vacuum is magnetic field free.

3.3 neutrino flavors and solar neutrinos

From experimental results[23] we can be sure that there are at least three kinds of flavors – ν_e, ν_μ, ν_τ . Up to now, it seems that we have assumed only one kind of neutrino because we cannot distinguish neutrino flavors with one longitudinal vacuum string oscillation model. Yet, there is a way to distinguish them. If neutrino propagation is assumed as a bundle of vacuum string oscillations in general, each flavor of neutrinos can be distinguished as a bulk motion with a different number of vacuum string oscillations.

If there exist the counter parts of left-handed neutrino ν_L and right-handed antineutrino $\bar{\nu}_R$ as assumed before, the parity should be conserved. Moreover, we found in the model of $W^\pm$ bosons that the helicity of neutrino is affected by the external magnetic field.

With these facts, let us try to find a possible explanation for the solar neutrino problem as mentioned before. For example, ^8B -neutrino deficit[28] as $\frac{\Phi_{exp.}}{\Phi_{SSM}} \approx 0.47 \pm 0.10(1\sigma)$ was confirmed again by Super-Kamiokande[26] experimental result. If solar magnetic field doesn't affect on helicities of neutrinos, the half of the neutrinos are *right handed*; the other half, *left handed*. Now, if we compare the cross-sections(leptonic interaction) of *right handed* neutrino and *left handed* neutrino to electron, the ratio of $\sigma(\bar{\nu}_R e^-)$ to $\sigma(\nu_L e^-)$ is ~ 0.416 for ^8B -neutrinos.[25] Hence, with this fact we can estimate that $\Phi_{exp.}$ should be $\sim 3.65 \times 10^6 \text{cm}^{-2} \text{s}^{-1}$. In fact, it is still bigger than the experimental result of $2.42 \pm 0.06 \times 10^6 \text{cm}^{-2} \text{s}^{-1}$.

Although it is not easy to estimate how much solar magnetic field affect on solar neutrinos(helicity and radiational direction), we can suppose that the neutrino flux itself from the Sun is not *isotropic*. In the model of W^+ boson in Fig.(2), we considered that virtual positive-charge-string is affected during the disintegration to e^+ and ν_e ($\sim 10^{-24} \text{sec}$) – Faraday Induction law. If neutrinos, those of which we are looking for on earth, emit from the equatorial zone of the Sun, then the direction of emission possibly can be deviated from the equatorial plane of the Sun. This effect should reduce the neutrino flux to 66%. Moreover, we can expect that the smaller energy neutrinos, the more strong deviations. In Kamiokande experimental result(1994)[26], we can confirm this effect by comparing neutrino spectrums in the measurements and from the Monte Carlo simulation, that is the expectation from standard solar model.

4 Summary

It has been known that Dirac equation represents spin $\frac{1}{2}\hbar$ fermions. Through this letter we compared Dirac formalism and Majorana formalism for neutrino case. Before that, we investigated how Pauli exclusion principle is related to fermions in the Complex Space[5] and if neutrino oscillation is physically feasible or not. In short, neutrino oscillation is not compatible with Special Theory of Relativity. That is, as long as Special Theory of Relativity is impeccable, neutrino flavor oscillation is not possible.

For neutrino, Majorana type neutrino was rejected in considering experimental results[16][17]. If neutrino has rest mass, it should be treated like other spin $\frac{1}{2}\hbar$ fermions. However, we assumed that neutrino has no mass; right-handed neutrino ν_R and left-handed antineutrino $\bar{\nu}_L$ exist. Here, we use neutrino and antineutrino, but they are not different even though we use them traditionally in Dirac four spinor formalism. There are only two physically distinctive neutrinos for each kind – *right-handed* and *left-handed*. From this fact and under the assumption that neutrino has no rest mass, the longitudinal vacuum string oscillation was suggested for neutrino. Moreover, the intermediate bosons $W^\pm$ was also suggested as a model because it is necessary that *parity* should be conserved if right-handed neutrino ν_R and left-handed antineutrino $\bar{\nu}_L$ exist. In the model of $W^\pm$ bosons, we also found how we might have overlooked the truth.

Meanwhile, V-A theory, with which the interaction Hamiltonian in Weak Interaction has been formulated, is not enough to include ν_R and $\bar{\nu}_L$. Instead, both V-A and V+A theories should be considered because we are assuming *parity* conservation – that means existing the mirror image. Moreover, the *lepton number* conservation law should be extended to include the other part, that might be a pair of *lepton number* conservations, separately.

The neutrino oscillation has been considered as a candidate to a new physics to explain the discrepancy of standard solar model[28] in solar neutrino experiments.[24][27] In spite of that, we investigated *neutrino* itself alternatively and suggested one way to solve the solar neutrino problem.

References

- [1] C. S. Wu *et al.*, *Phys. Rev.* **105**, 1413 (1957) ; Richard L. Garwin, Leon M. Lederman, Marcel Weinrich, *Phys. Rev.* **105**, 1415(1957) ; G. Backenstoss *et al.*, *Phys. Rev.Lett.* **6**, 8, 415(1961) ; Marcel Bardon, Paolo Franzini, Juliet Lee, *Phys. Rev.Lett.* **7**, 1, 23(1961)
- [2] Donald H. Perkins, *Introduction to High Energy Physics*, 3rd edition, (Addison-Wesley publishing Company, Inc., 1987), pp. 23
- [3] Hans Fraunfelder, Ernest M. Henley, *Subatomic Physics* , (Prentice-Hall, Inc., 1974), pp. 79, Chap 10.8 ; D.O. Caldwell, *et al*, *Phys. Rev.* **D**, 7, 1362(1973) ; *Phys. Rev. Lett.* **25**, 609, 613(1970) ; L. Stodolsky, *Phys. Rev. Lett.* **18**, 135 (1967)
- [4] W. Greiner, B. Müller, *Gauge Theory of Weak Interaction*, (Springer-Verlag Berlin Heidelberg, 1996), pp. 1.
- [5] Kiyoun Kim, *quant-ph/9701023*, e-print archive
- [6] Kiyoun Kim, *quant-ph/9706063*, e-print archive
- [7] P.A.M Dirac, *Proc. Roy. Soc. (London)* **A 126**, 360 (1930)
- [8] Berry, M.V., *Proc. Roy. Soc.(London)* **A 392**, 45 (1984)
- [9] Thomas F. Jordan, *J. Math. Phys.* **29**(9), 2042 (1988)
- [10] E. Majorana, *Nuovo Cimento* **14**, 171 (1937) ; Claude Itzykson, Jean-Bernard Zuber, *Quantum Field Theory*, (McGraw-Hill Inc., 1980), pp. 49-50, 164
- [11] K. M. Case, *Phys. Rev.* **107**, 307 (1957)
- [12] F. Boehm, P. Vogel, *Physics of Massive Neutrino 2nd edition*, (Cambridge University Press, Cambridge, 1992), pp. 1-20
- [13] Claude Itzykson, Jean-Bernard Zuber, *Quantum Field Theory*, (McGraw-Hill Inc., 1980), pp. 45-104
- [14] Lewis H. Ryder, *Quantum Field Theory*, (Cambridge University Press, 1985), pp. 31-57
- [15] Claude Itzykson, Jean-Bernard Zuber, *Quantum Field Theory*, (McGraw-Hill Inc., 1980), pp. 85-89
- [16] Cowan Jr. and F. Reines, *Phys. Rev.* **106**, 825 (1957) ; R. Davis, *Phys. Rev.* **97**, 766 (1953)

- [17] L.A. Ahrens *et al*, *Phys. Rev. Lett.* **54**, 18(1985) ; Vernon D. Barger, Roger J. N. Phillips, *Collider Physics*, (Addison-Wesley publishing Co., 1987), pp. 133 ; Klaus Winter *Neutrino Physics*, (Cambridge University Press, 1991) pp. 613
- [18] T.D. Lee, C.N. Yang, *Phys. Rev.* **105**, 5, 1671 (1957)
- [19] R.J. Protheroe *et al*, *AP. J. Lett.* 280,L47(1984) ; R.M. Baltrusaitis *et al*, *AP. J. Lett.* 293, L69(1985) ; R.J. Protheroe and R.W. Clay, *Nature* **315**, 205,(1985)
- [20] John N. Bahcall,*et al.*, *Physical Review D* **51**, 6146(1995) ; Vernon D. Barger, Roger J. N. Phillips, *Collider Physics*, (Addison-Wesley publishing Co., 1987), pp. 133 ; Francis Halzen, Alan D. Martin, *Quark and Leptons*, (John Wiley & Sons, Inc., 1984), pp. 267-271 ; Claude Itzykson, Jean-Bernard Zuber, *Quantum Field Theory*, (McGraw-Hill Inc., 1980), pp. 224-230
- [21] J.J. Sakurai, *Advance Quantum Mechanics*, (Addison-Wesley Publishing Company, Inc., 1980) pp. 138-139 ; James D. Bjorken, Sidney D. Drell, *Relativistic Quantum Mechanics*, (McGraw-Hill, Inc., 1964) pp. 70-71
- [22] Gordon Kane, *Modern Elementary Particle Physics*, (Addison-Wesley Publishing Company, 1987), pp. 122-128
- [23] M.L. Perl *et al.*, *Phys. Rev. Lett.* **35** , 1489 (1975) ; G. Dandy *et al.*, *Phys. Rev. Lett.* **9**, 36 (1962) ; Leon M. Lederman, *Sci. Amer.* **208**, 3, 60(1963)
- [24] William J. Marciano, *Comments Nucl. Part. Phys.*, 1981, Vol. 9, No. 5, pp. 169-182
- [25] Klaus Winter *Neutrino Physics*, (Cambridge University Press, 1991) pp. 586-597, 613
- [26] The Kamiokande Collaboration, *Phys. Lett.* **B**, 335, 237(1994) ; The Super-Kamiokande Collaboration, *hep-ex/9805021*, e-print archive
- [27] Chritine Sutton, *Spaceship Neutrino*, (Cambridge University Press, 1992), pp. 148-183
- [28] John N. Bahcall, *Phys. Lett.* **B**, 433, 1(1998) ; Solar Fusion Cross Sections *astro-ph/9805121*, e-print archive